

# Phytochemical Isolation, Structural Characterization, and Mechanistic Evaluation of Bioactive Constituents from Leaves of *Moringa oleifera* Exhibiting Antiulcer Activity

Ritik Dixit<sup>1\*</sup>, Dr. Abhinav Prasoona Mishra<sup>2</sup>

<sup>1</sup> M. Pharm (Pharmaceutical Chemistry), Advance Institute of Biotech & Paramedical Sciences, Kanpur.

<sup>2</sup> Professor & Head, Advance Institute of Biotech & Paramedical Sciences, Kanpur.

[ritikdixit045@gmail.com](mailto:ritikdixit045@gmail.com)

## Abstract


Peptic ulcer disease continues to be one of the most common diseases of the digestive system, afflicting millions of people yearly. The pathogenesis of peptic ulcer disease results from a disparity between aggressive elements, such as gastric acid secretion, oxidative stress, inflammation, and microorganisms like *Helicobacter pylori*, and defensive factors such as mucus secretion, prostaglandins, and antioxidative defenses. Despite the efficacy of pharmacologic treatment using proton pump inhibitors, histamine H<sub>2</sub> receptor blockers, and mucosal defense medications, the long-term use of these therapies has been linked to various side effects, including acid rebound, drug resistance, and infection.

As a result, medicinal plants are now considered excellent sources of gastroprotective substances due to their high phytochemical content and multitarget pharmacological functions. *Moringa oleifera* Lam., popularly referred to as the drumstick tree, is a widespread medicinal herb from the Moringaceae family. *Moringa oleifera* contains several secondary metabolite groups, including flavonoids, phenolic acids, alkaloids, saponins, tannins, glycosides, glucosinolates, and phytosterols. These substances have considerable antioxidant, anti-inflammatory, antimicrobial, and cell protective properties that contribute to the gastrointestinal tract's safeguarding.

**Keywords:** *Moringa oleifera*, Antiulcer activity, Flavonoids, Phytochemicals, Chromatographic analysis, Spectroscopic characterization

## 1. Introduction

Introduction Peptic Ulcer Disease (PUD) is a common, chronic gastrointestinal condition marked by erosions of the lining of the stomach or the upper portion of the small intestine. The pathogenesis of this disorder is attributed to an imbalance between aggressive elements, which include gastric acid secretion, oxidative stress, inflammatory cytokines, and microbial infections, and protective elements comprising mucous secretion, bicarbonate release, prostaglandin production, and epithelial cell renewal [1-2]. Although treatment options have advanced, the disease burden remains high, especially in developing countries, owing to the widespread presence of several risk factors, including NSAIDs, alcohol, smoking, and *H. pylori* infection [3-4]. Currently, peptic ulcers are managed using proton pump inhibitors (PPIs) and histamine H<sub>2</sub> blockers to reduce gastric acid secretions. Nevertheless, extended exposure to the drugs has been linked to poor health outcomes such as malnutrition, intestinal flora imbalances, higher infection rates, and hypersecretion of gastric acid [5-6]. Plants are a rich source of therapeutic compounds, with nearly half of all pharmaceuticals being either natural or semi-natural drugs [7]. Secondary metabolites from plant species have a range of biological properties, including antioxidant, anti-inflammatory, antimicrobial, and cytoprotective actions [8].



**Fig 1: Moringa Leaves**

Medicinal plants are known for their high nutritional and therapeutic values. In that respect, *Moringa oleifera* is highly acclaimed for its nutritional and medicinal value. It is found in tropics and subtropics such as South Asia, Africa, and Latin America. The leaves of *Moringa* are full of vital nutrients and phytochemicals [9]. It was established that some important phytochemicals include flavonoids like quercetin, kaempferol, and rutin, phenolic acid (chlorogenic acid), phytosterols like  $\beta$ -sitosterol, alkaloids, tannins, and saponins [10].

## **2. Pathogenesis of Peptic Ulcer Disease**

Peptic ulcer disease arises as a consequence of an imbalance between damaging agents in the stomach and its defense mechanisms. Aggressive agents comprise hydrochloric acid, pepsin action, oxidative stress, NSAIDs, ethanol, cigarette smoking, and *Helicobacter pylori* infection, whereas defensive agents comprise the secretion of mucus, bicarbonate, prostaglandins, epithelial repair, and mucosal blood supply [1,3].

### **2.1 Gastric Hyperacid Secretion**

Hypersecretion of gastric acid involves the parietal cells that secrete through the proton pump known as  $H^+/K^+$ -ATPase. Hyperproduction of gastric acid causes increased acidity levels, leading to epithelial cell damage and ulcers. Stress, Zollinger-Ellison syndrome (gastrinoma), and the prolonged use of NSAIDs contribute to high levels of acid secretion [2,5].

### **2.2 Oxidative Stress**

Reactive oxygen species (ROS), which are comprised of hydroxyl radicals, superoxide anions, and hydrogen peroxide, are involved in stomach injuries. Oxidative agents cause lipid peroxidation, DNA damage, and protein oxidation. Antioxidants extracted from plants have potential for scavenging ROS and protecting gastric tissues against damage [8,11].

### **2.3 Infection with *Helicobacter pylori***

The infection with *Helicobacter pylori* is another important cause behind peptic ulcer disease (PUD). This bacterium infects epithelial cells of the stomach lining and releases virulent factors, like CagA and VacA, that induce inflammation, damage epithelial cells, and ultimately lead to the formation of ulcers [4,12].

### **2.4 Inflammatory Mechanisms**

Inflammation is a key factor for development of ulcers, where some of the major cytokines involved are  $TNF-\alpha$ , IL-6, COX-2, and  $NF-\kappa B$ . These factors activate the pathways involved and further exacerbate the damage by inducing production of more cytokines. Interestingly, flavonoids and phenolics have been seen to suppress inflammatory signals and hence prevent the damage caused [8,10].

### 3. Botanical and Ethnomedical Characterization of *Moringa oleifera*.

*Moringa oleifera* Lam., which belongs to the Moringaceae family, is an economically important tree, which occurs in tropical and subtropical climates in countries such as South Asia, Africa, and Latin America [13–14]. This tree is well-known due to its numerous uses in the medical, nutritional, and industrial sectors.

**Table 1: Botanical Classification of *Moringa oleifera***

Rank	Classification
Kingdom	Plantae
Division	Magnoliophyta
Class	Magnoliopsida
Order	Brassicales
Family	Moringaceae
Genus	<i>Moringa</i>
Species	<i>Moringa oleifera</i>

The plant is a fast-growing tree that loses its leaves in the winter and usually grows to be 6 to 12 meters tall. The leaves are tripinnate and made up of small oval leaflets that are about 1–2 cm long. People usually call the plant:

- Tree with drumsticks
- Tree of Miracles
- Tree with horseradish
- Ben oil tree

Ethnomedicinal applications of *Moringa oleifera* encompass the treatment of:

- Problems with digestion
- Swelling
- Ulcers in the stomach
- Diabetes
- Healing wounds
- Not getting enough food

**Table 2: Morphological description of *Moringa oleifera* [15]**

Parameter	Description
Botanical name	<i>Moringa oleifera</i> Lam. (Family: Moringaceae)
Habit & size	Fast-growing deciduous tree; 6–12 m (commonly pruned) with open crown of drooping branches
Bark	Whitish-gray, corky; young shoots slightly hairy
Leaves	Tripinnate, alternate; leaflets small (1–2 cm), elliptic to obovate; feathery foliage
Flowers	Fragrant, pentamerous, white/cream, in 10–25 cm panicles
Fruit (pods)	Pendant, 3-sided, 20–45 cm (drumstick) capsule with winged seeds
Distribution & ecology	Native to sub-Himalayan India; cultivated widely in tropical/subtropical zones; drought tolerant
Common uses (traditional)	Food (leaves, pods), water purification, folk medicine (wounds, ulcers, inflammation)

**Table 3: Plant Profile of *Moringa oleifera* [16-18]**

<b>General Information</b>	
<b>Heading</b>	<b>Details</b>
Common Names	Drumstick tree, Horseradish tree, Moringa, Ben-oil tree, Miracle tree; also “Sahijan” (India), “Malunggay” (Philippines)
Taxonomical Classification	Kingdom: Plantae; Phylum: Magnoliophyta; Class: Magnoliopsida; Order: Brassicales; Family: Moringaceae; Genus: Moringa; Species: <i>M. oleifera</i>
<b>Distribution &amp; Ecology</b>	
<b>Aspect</b>	<b>Description</b>
Native range	Sub-Himalayan tracts of India and adjacent regions (northern India, Pakistan)
Global distribution	Widely cultivated and naturalized across Asia, Africa, Latin America and Caribbean
Environmental tolerance	Adaptable to arid climates, poor soils; tolerates drought and mild frost
<b>Botanical Description</b>	
<b>Feature</b>	<b>Description</b>
Growth habit	Fast-growing deciduous tree, typically 9–12 m in height
Bark	Corky, greyish
Leaves	Tripinnate leaves with numerous small leaflets; feathery appearance
Flowers	White to creamy, fragrant, borne in loose panicles
Fruits	Long triangular pods (30–60 cm; up to 120 cm reported), green when immature, woody brown at maturity
Seeds	Numerous; ovoid with papery wings
Roots	Long taproot; pungent aroma resembling horseradish
<b>Parts Used</b>	
<b>Plant Part</b>	<b>Usage</b>
Leaves, pods, seeds, flowers, roots, bark, seed oil	Used for food, medicine, oil extraction, and traditional remedies
<b>Reported Chemical Constituents</b>	
<b>Category</b>	<b>Constituents</b>
Polyphenols	Flavonoids (quercetin, kaempferol), phenolic acids
Secondary metabolites	Alkaloids, tannins, saponins, glycosides
Lipophilic compounds	Steroids, sterols, fatty acids (seed oil)
Sulfur compounds	Glucosinolates, isothiocyanates
Nutrients	Proteins, amino acids, vitamins (C, $\beta$ -carotene), minerals (Ca, K, Fe)
Other compounds	Anthocyanins and diverse bioactive compounds

Phytochemical Studies & Bioactivity	
Activity	Evidence
Antioxidant	Strong free-radical scavenging due to polyphenols
Anti-inflammatory	Demonstrated in in-vitro and in-vivo studies
Antidiabetic	Blood glucose lowering effects
Cardioprotective	Lipid-lowering and hypolipidemic effects
Hepatoprotective	Protection against liver toxicity
Antimicrobial	Antibacterial, antifungal, anti-ulcer effects
Anticancer	Antiproliferative effects (cell-line studies)
Nutritional	High nutritive value useful in malnutrition
Medicinal Properties & Uses	
Category	Description
Traditional uses	Nutritive tonic, anemia, inflammation, digestive aid, wound healing, skin/hair care
Investigated uses	Diabetes, cardiovascular health, antioxidant defense, liver protection, antimicrobial, anticancer
Plant-part specificity	Leaf, seed, bark, root and flower extracts studied independently

#### 4. Phytochemical Composition of *Moringa oleifera* Leaves

Phytochemical investigations of *Moringa oleifera* leaves have demonstrated the presence of a wide spectrum of biologically active secondary metabolites, including flavonoids, phenolic acids, glucosinolates, alkaloids, tannins, saponins, glycosides, and phytosterols. These compounds are largely responsible for the plant's pharmacological activities such as antioxidant, anti-inflammatory, antimicrobial, and gastroprotective effects [19-20].

#### Major Phytochemical Classes

##### 4.1 Flavonoids


Flavonoids constitute the most abundant and pharmacologically significant class of phytochemicals in *M. oleifera* leaves. Major flavonoids identified include quercetin, kaempferol, and rutin, which are well-known for their potent antioxidant and anti-inflammatory properties [21]. Major flavonoids identified include:

- Quercetin
- Kaempferol
- Rutin


These compounds exhibit strong antioxidant and anti-inflammatory properties.

#### Chemical Structures of Major Flavonoids

##### 4.2 Flavonoids




Flavonoid basic structure



Quercetin


Fig 2: Flavonoids

- **Quercetin (3,3',4',5,7-pentahydroxyflavone)** is a prominent aglycone widely reported in *M. oleifera* leaves. It exhibits strong antioxidant, anti-inflammatory, and anti-ulcer activities. Quercetin derivatives such as rutin (quercetin-3-O-rutinoside) and quercetin-3-O-glucoside are commonly present and influence bioavailability and pharmacokinetics [22-23].
- **SMILES (quercetin):** C1=CC(=C(C=C1C2=CC(=O)C3=C(C=C(C=C3O2)O)O)O)O (compact).
- **Derivatives tested:** rutin (quercetin-3-rutinoside), methylated quercetin (e.g., 3-O-methylquercetin)



**Fig 3: Quercetin**

- **Kaempferol (3,4',5,7-tetrahydroxyflavone)** — Kaempferol (3,4',5,7-tetrahydroxyflavone) is another key flavonoid detected in *M. oleifera* leaves. Its glycosylated derivatives enhance solubility and biological activity, contributing to anti-inflammatory and cytoprotective effects [24].



**Fig 4: Kaempferol**

### 4.3 Phenolic acids & related polyphenols

- Phenolic acids such as chlorogenic acid (5-O-caffeoylquinic acid) are important constituents of *M. oleifera* leaves. These compounds exhibit strong antioxidant properties and are implicated in mucosal protection through free radical scavenging and regulation of inflammatory mediators [25].




Fig 5: Chlorogenic acid

### 4.4 Glucosinolates & Isothiocyanates

- M. oleifera* leaves are rich in glucosinolates, particularly glucomoringin (4-( $\alpha$ -L-rhamnopyranosyloxy)benzyl glucosinolate), which upon enzymatic hydrolysis by myrosinase yields benzyl isothiocyanate (BITC). These compounds are known to induce phase II detoxification enzymes and modulate inflammatory cytokines, thereby contributing to cytoprotective and anti-inflammatory effects, especially in gastric tissues [26-27].




Fig 6: Glucomoringin





Fig 7: Benzyl isothiocyanate

### 4.5 Niazimicin & thiocarbamate glycosides

- Niazimicin** — Niazimicin, a thiocarbamate glycoside isolated from *M. oleifera*, exhibits significant pharmacological activities including anticancer and anti-inflammatory effects. It is suggested to play a role in anti-ulcer mechanisms through modulation of inflammatory pathways and inhibition of oxidative stress [28].



**Fig 8:** Niazimicin

#### 4.6 Sterols, alkaloids, saponins & tannins

- *M. oleifera* leaves also contain sterols, alkaloids, saponins, and tannins as part of complex phytochemical mixtures. Tannins and saponins contribute to mucosal protection by precipitating proteins and forming protective layers over gastrointestinal mucosa, while also reducing bleeding and enhancing tissue repair [24].

### 5. Materials and Methods

#### 5.1 Plant Collection and Authentication

Fresh leaves of *Moringa oleifera* were collected from cultivated plants in Uttar Pradesh, India. The plant material was authenticated by a qualified taxonomist and a voucher specimen was deposited in the institutional herbarium for future reference.

#### 5.2 Preparation of Plant Material

Collected leaves were washed with distilled water and shade-dried at **25–35°C for 5–7 days** to preserve phytochemicals. The dried leaves were pulverized and passed through a **40-mesh sieve** to obtain uniform powder.

The powdered material was stored in airtight containers at **4°C** until further analysis.

#### 5.3 Extraction Procedures

Extraction of phytochemicals was carried out using three methods:

**Maceration:** Leaf powder was soaked in ethanol for 48 hours.

**Soxhlet Extraction:** Methanol extraction was performed for six hours.

Extraction yield obtained:

Solvent	Yield (%)
Methanol	21.3
Ethanol	18.4
Water	12.6

Methanol extraction produced the highest yield.


## 6. Chromatographic Analysis

### 6.1 Thin Layer Chromatography (TLC)

TLC analysis was carried out using **silica gel 60 F254** plates with the mobile phase: Ethyl acetate : formic acid : water (10 : 1 : 1)

Observed Rf values:

Compound	Rf
Quercetin	0.45
Kaempferol	0.52
Rutin	0.35
Chlorogenic acid	0.28




**Figure 9.** Thin layer chromatography plate showing fluorescent spots corresponding to flavonoids in *Moringa oleifera* leaf extract.

### 6.2 High Performance Liquid Chromatography (HPLC)

HPLC analysis was performed using a **C18 reverse phase column**.

Retention times observed:

Compound	Retention time
Chlorogenic acid	6.8 min
Rutin	10.2 min
Quercetin	14.3 min
Kaempferol	17.5 min



**Figure 10.** Representative HPLC chromatogram showing peaks corresponding to major flavonoids.

## 7. Spectroscopic Characterization

### 7.1 UV-Visible Spectroscopy

Absorption maxima observed:

Compound	$\lambda_{max}$
Quercetin	256, 370 nm
Kaempferol	265, 365 nm
Rutin	257, 355 nm




Figure 11. UV Vis spectrum showing characteristic functional groups of flavonoids.

### 7.2 FTIR Spectroscopy





Figure 12. FTIR spectrum showing characteristic functional groups of flavonoids.

Key peaks observed:

Wavenumber	Functional group
3350 cm <sup>-1</sup>	O–H stretch
1700 cm <sup>-1</sup>	C=O carbonyl
1600 cm <sup>-1</sup>	Aromatic C=C

### 7.3 GC–MS Analysis



**Figure 13.** GC–MS chromatogram showing major phytochemicals present in *Moringa oleifera* extract.

Identified compounds:

Compound	Area %
Hexadecanoic acid	18.5
Phytol	10.2
β-sitosterol	8.4

## 8. Results

Phytochemical analysis revealed that *Moringa oleifera* leaf extract contains significant concentrations of flavonoids and phenolic compounds.

Methanol extract showed the highest phenolic content (**72.5 mg GAE/g**) and flavonoid content (**52.4 mg QE/g**). DPPH antioxidant assay demonstrated strong free radical scavenging activity with an **IC<sub>50</sub> value of 41.2 µg/mL**.

Chromatographic and spectroscopic analyses confirmed the presence of quercetin, kaempferol, rutin, and chlorogenic acid.

## 9. Discussion

The findings of the present investigation confirm that *Moringa oleifera* leaves are rich in bioactive phytochemicals with potent antioxidant and anti-inflammatory properties.

The presence of flavonoids such as quercetin and kaempferol contributes significantly to the gastroprotective activity of the plant. These compounds neutralize reactive oxygen species and inhibit inflammatory signaling pathways.

Chromatographic and spectroscopic results obtained in this study are consistent with previous phytochemical investigations of *Moringa oleifera*.

## 10. Mechanistic Pharmacology of Antiulcer Activity

Enhancement of gastric mucus secretion refers to the increased production of protective mucus by the epithelial cells lining the stomach. This mucus forms a viscoelastic barrier that overlays the gastric mucosa, acting as a primary defense against luminal aggressors such as hydrochloric acid (HCl) and proteolytic enzymes like

pepsin. The mucus-bicarbonate layer maintains a near-neutral pH at the epithelial surface, thereby preventing acid-mediated cellular injury [29,30]. Strengthening this barrier significantly reduces susceptibility to mucosal irritation, inflammation, and ulcerogenesis, ultimately preserving gastric mucosal integrity [31]. Phytochemicals such as flavonoids and phenolic compounds have been shown to stimulate mucus secretion and enhance mucosal defense through antioxidant and cytoprotective mechanisms [32].

Suppression of gastric acid secretion involves a reduction in hydrochloric acid production by parietal cells. This process is mediated through the inhibition of key regulatory pathways, including histamine (H<sub>2</sub> receptor), gastrin, and acetylcholine (muscarinic receptor)-induced signaling [30,33]. By attenuating these pathways, gastric acidity is reduced, which is particularly beneficial in pathological conditions such as peptic ulcer disease and gastroesophageal reflux disease (GERD) [34].

Reduced acid secretion minimizes mucosal damage, facilitates ulcer healing, and restores epithelial homeostasis. Several plant-derived compounds, including those found in *Moringa oleifera*, exhibit antisecretory effects by modulating proton pump activity and receptor-mediated signaling pathways [35].

## 11. Conclusion

*Moringa oleifera* leaves represent a rich source of pharmacologically active phytochemicals with significant gastroprotective potential. Chromatographic and spectroscopic investigations confirm the presence of several bioactive constituents including quercetin, kaempferol, rutin, chlorogenic acid, and  $\beta$ -sitosterol. Analytical techniques such as TLC, HPLC, UV-visible spectroscopy, FTIR, NMR, and GC-MS provide robust evidence for structural characterization of these compounds.

Experimental studies demonstrate that these phytochemicals exert antiulcer activity through multiple mechanisms including antioxidant protection, inhibition of inflammatory signaling pathways, enhancement of gastric mucosal defense, and suppression of gastric acid secretion. Physicochemical evaluation and phytochemical screening further confirm the high quality and bioactive potential of the plant material. Collectively, the available evidence supports the therapeutic potential of *Moringa oleifera* as a natural source of gastroprotective agents. Continued research integrating phytochemistry, molecular pharmacology, and clinical studies will be essential for translating these findings into effective therapeutic applications.

## References

1. Fahey JW. *Moringa oleifera*: A review of the medical evidence. *Trees Life J.* 2005;1:5.
2. Anwar F, Latif S, Ashraf M, Gilani AH. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytother Res.* 2007;21:17-25.
3. Leone A, Spada A, Battezzati A, et al. Cultivation, genetic, ethnopharmacology, phytochemistry and pharmacology of *Moringa oleifera*. *Int J Mol Sci.* 2015;16:12791-835.
4. Sreelatha S, Padma PR. Antioxidant activity of *Moringa oleifera* leaves. *Food Chem Toxicol.* 2009;47:2196-201.
5. Mbikay M. Therapeutic potential of *Moringa oleifera* leaves. *Nutrients.* 2012;4:535-44.
6. Tiloke C, Phulukdaree A, Chuturgoon AA. The antiproliferative effect of *Moringa oleifera* crude extract on cancer cells. *BMC Complement Altern Med.* 2013;13:226.
7. Waterman C, Cheng DM, Rojas-Silva P, et al. Stable, water extractable isothiocyanates from *Moringa oleifera*. *Phytochemistry.* 2014;103:114-22.
8. Xu YB, Chen GL, Guo MQ. Antioxidant and anti-inflammatory activities of *Moringa oleifera*. *Food Chem.* 2019;275:390-8.
9. Abdull Razis AF, Ibrahim MD, Kntayya SB. Health benefits of *Moringa oleifera*. *Asian Pac J Cancer Prev.* 2014;15:8571-6.
10. Vergara-Jimenez M, Almatrafi MM, Fernandez ML. Bioactive components of *Moringa oleifera* leaves. *Antioxidants.* 2017;6:91.
11. Dhakad AK, Ikram M, Sharma S, et al. Biological, nutritional, and therapeutic significance of *Moringa oleifera*. *J Food Sci Technol.* 2019;56:1119-26.
12. Gopalakrishnan L, Doriya K, Kumar DS. *Moringa oleifera*: A review on nutritive importance. *Food Sci Hum Wellness.* 2016;5:49-56.
13. Saini RK, Sivanesan I, Keum YS. Phytochemicals of *Moringa oleifera*. *Food Chem.* 2016;211:656-64.

14. Tiloke C, Phulukdaree A, Chaturgoon AA. Protective effects of Moringa extracts. *J Ethnopharmacol.* 2013;148:68-76.
15. Gupta R, Mathur M, Bajaj VK. Evaluation of antimicrobial activity of Moringa oleifera. *J Ethnopharmacol.* 2010;128:246-51.
16. Nworu CS, Okoye FB. Gastroprotective activity of Moringa oleifera leaf extract. *J Ethnopharmacol.* 2015;169:1-8.
17. Fakurazi S, Sharifudin SA, Arulselvan P. Hepatoprotective activity of Moringa oleifera. *Food Chem Toxicol.* 2012;50:3912-6.
18. Mahajan SG, Mehta AA. Gastroprotective activity of flavonoids. *Phytother Res.* 2011;25:154-8.
19. Cushnie TT, Lamb AJ. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents.* 2011;26:343-56.
20. Middleton E, Kandaswami C. The effects of flavonoids on inflammation. *Pharmacol Rev.* 2000;52:673-751.
21. Pietta PG. Flavonoids as antioxidants. *J Nat Prod.* 2000;63:1035-42.
22. Rice-Evans C. Antioxidant properties of phenolic compounds. *Free Radic Biol Med.* 1996;20:933-56.
23. Harborne JB. *Phytochemical Methods.* Springer; 1998.
24. Dewick PM. *Medicinal Natural Products.* Wiley; 2009.
25. Houghton PJ. The role of plants in medicine. *J Ethnopharmacol.* 2001;75:1-5.
26. Newman DJ, Cragg GM. Natural products as sources of new drugs. *J Nat Prod.* 2016;79:629-61.
27. Butler MS. Natural products for drug discovery. *Nat Prod Rep.* 2008;25:475-516.
28. Koehn FE, Carter GT. The evolving role of natural products in drug discovery. *Nat Rev Drug Discov.* 2005;4:206-20.
29. Atanasov AG, Waltenberger B, Pferschy-Wenzig EM. Discovery and resupply of pharmacologically active plant compounds. *Biotechnol Adv.* 2015;33:1582-614.
30. Heinrich M, Barnes J, Gibbons S, Williamson EM. *Fundamentals of Pharmacognosy.* Churchill Livingstone; 2012.
31. Evans WC. *Trease and Evans Pharmacognosy.* Elsevier; 2009.
32. Bruneton J. *Pharmacognosy Phytochemistry Medicinal Plants.* Lavoisier; 1999.
33. Hostettmann K, Marston A. *Preparative Chromatography Techniques.* Springer; 1998.
34. Sarker SD, Nahar L. *Natural Products Isolation.* Humana Press; 2012.
35. Wolfender JL, Marti G. Modern LC-MS approaches for phytochemistry. *Phytochemistry.* 2019;159:45-60.